

Sever's disease

What is sever's disease?

Sever's disease causes pain in the bone at the back of the heel. This pain occurs because of a mismatch between the growth of the calf bones and muscles.

What are the symptoms of Sever's disease?

Typically the pain at the back of the heel is worse after physical activity and improves with rest. Sever's disease can affect one or both feet.

Sever's disease is slightly more common in boys than girls and usually affects children aged 8 – 12 years. It is uncommon in older teenagers as the heel bone usually finishes growing by age 15 years.

What causes Sever's disease?

During a growth spurt, the bones grow first and the muscles and tendons have to catch up. The Achilles tendon is a large tendon which attaches the muscles of the calf to the back of the heel. In Sever's disease, the heel bone becomes sore and swollen where the Achilles tendon attaches to it.

Children who are involved in running and jumping sports are those who tend to get Sever's disease most often. These include basketball, soccer, Australian Rules football and some forms of athletics. Wearing shoes with stops (studs) increases the risk of developing Sever's disease.

How is it diagnosed?

Your child's doctor will diagnose Sever's disease from the typical symptoms and appearance. There is rarely a need for blood tests or x-rays.

How is Sever's disease treated?

Your local doctor or physiotherapist can give you advice about managing Sever's disease. Specialist referral is only required for very unusual cases.

Rest and cold packs on the painful area can help during flare ups. Cushion pads in the heels of shoes (available from pharmacists, podiatrists and physiotherapists) may help. Stretching exercises of the Achilles tendons are important. A physiotherapist can teach these to you and your child.

Generally, medication is not recommended. Paracetamol (e.g. Panadol) or anti-inflammatory medicines such as ibuprofen (e.g. Nurofen) may be used for short periods of bad pain.

There is no evidence that complementary medicines, including acupuncture, are helpful for Sever's disease.

How can I help my child?

Reassure your child that the pain will go away in time. Seeing a physiotherapist can be helpful to learn ways to stretch the Achilles tendon and keep pain under control. Encourage your child to use other strategies, such as rest and cold packs, to manage their pain.

Living with Sever's disease

Sever's disease will go away with time as growth of the heel bone finishes. In the meantime, pain may be worse with activity but there are things that you and your child can do to make life easier.

Changes in sporting activities may help to reduce pain. Fewer running and jumping activities will result in less pain. Continuing to do these activities will not damage your child's foot, but may make the pain worse.

ARTHRITIS

INFORMATION SHEET


Increasing other activities may be important for your child in the meantime, such as swimming or cycling. It may be helpful for coaches and sports teachers to be aware of your child's condition. Reducing the use of shoes with studs can make a big difference.

What is the outlook for my child?

The pain of Sever's disease will disappear with time but may come back at times. Sever's disease often lasts two to three years. Once your child has finished their growth spurt the pain will go away. Your child will not have any long term problems because of this condition.

CONTACT YOUR LOCAL ARTHRITIS OFFICE FOR MORE INFORMATION SHEETS ON ARTHRITIS.

Exercises that may help



i
reproduced with permission.

This sheet was produced in association with the Australian Paediatric Rheumatology Group

APRG
AUSTRALIAN PAEDIATRIC
RHEUMATOLOGY GROUP

© Copyright Arthritis Australia March 2015. Reviewed December 2017.
The Australian Paediatric Rheumatology Group contributed to the development of this information sheet.

Your local Arthritis Office has information, education and support for people with arthritis
Infoline 1800 011 041 www.arthritisaustralia.com.au

Disclaimer: This sheet is published by Arthritis Australia for information purposes only and should not be used in place of medical advice.